

Cemaliye Boylu Akyerli · Uğur Özbek
Müge Aydın-Sayitoğlu · Sema Sırma · Tayfun Özçelik

Analysis of *MYH* Tyr165Cys and Gly382Asp variants in childhood leukemias

Received: 4 March 2003 / Accepted: 4 July 2003 / Published online: 13 August 2003
© Springer-Verlag 2003

Introduction

DNA-repair gene mutations have been suspected as being a predisposing factor in the development of leukemia (Horwitz 1997). Our group identified one of the first examples of a DNA-repair gene mutation to be causally linked to childhood hematological malignancies and neurofibromatosis type I, which involves a homozygous germ-line mutation in the mismatch repair (MMR) gene *MLH1* (Ricciardone et al. 1999; Wang et al. 1999). Subsequently, homozygous inactivation of *MSH2*, another MMR gene, was also found to be associated with early onset leukemia (Whiteside et al. 2002). It is well known that heterozygous germ-line mutations in the MMR pathway genes *MLH1*, *MSH2*, *PMS2*, *PMS1*, and *MSH6* lead to hereditary non-polyposis colorectal cancer (HNPCC) (Peltomäki 2001). Thus, tumorigenesis through an “MMR deficiency pathway” appears to be associated with two different disease phenotypes which are dependent on the status of the germ-line mutation: (a) HNPCC when the mutation is present on only one allele (heterozygous), and (b) hematological malignancies when the mutation(s) is present on both alleles (homozygous or compound heterozygous).

Base excision repair (BER) is another important DNA-repair pathway and plays a significant role in the repair of mutations generated by reactive oxygen species during aerobic metabolism. BER was not linked with any human genetic disorder until recently, when a

British family in which three siblings affected by multiple colorectal adenomas and carcinoma was shown to be compound heterozygous for *MYH* missense variants Tyr165Cys (Y165C) and Gly382Asp (G382D) (Al-Tassan et al. 2002). *MYH* is a homologue of *E. coli* mutY, and the mutations mentioned above affect residues that are conserved (Tyr82 and Gly253). Tyrosine 82 is predicted to function in mismatch specificity and is located in the pseudo-helix-hairpin-helix (Guan et al. 1998). Adenine glycosylase activity assays of the Tyr82Cys and Gly253Asp mutant proteins with 8-oxoG:A and G:A substrates show that their rate for adenine removal at 37°C is reduced by approximately 98% (Tyr82Cys) and 86% (Gly253Asp) (Al-Tassan et al. 2002). Furthermore, bi-allelic germ-line mutations in *MYH* were identified in seven unrelated patients with colorectal adenomas (six with colorectal cancer) (Jones et al. 2002). Interestingly, the missense variations Tyr165Cys and Gly382Asp, which significantly reduce the adenine glycosylase activity of *MYH* protein, were each identified once in a normal control group of 100 British individuals with no history of colorectal adenoma or carcinoma (Al-Tassan et al. 2002). Since a connection between DNA-repair gene mutations and the path to hematological malignancy is now well established, and individuals who carry heterozygous *MYH* missense mutations Tyr165Cys and Gly382Asp have been documented in a control group (Al-Tassan et al. 2002), we investigated the association between these two *MYH* missense mutations and childhood leukemia risk.

C. B. Akyerli · T. Özçelik (✉)
Department of Molecular Biology and Genetics, Faculty
of Science, Bilkent University, Bilkent, 06800 Ankara, Turkey
E-mail: tozcelik@fen.bilkent.edu.tr

T. Özçelik
Ayhan Sahenk Foundation, 80670 Istanbul, Turkey

U. Özbek · M. Aydın-Sayitoğlu · S. Sırma
Department of Genetics, Institute for Experimental Medicine,
Istanbul University, Çapa, 34280 Istanbul, Turkey

Results and discussion

The study population included 185 cases of childhood leukemias subdivided into two groups: acute myeloid leukemia (AML; $n=45$) and acute lymphoblastic leukemia (ALL; $n=140$) diagnosed at Istanbul University between 1998 and 2002. Detailed clinical data are available for all patients. The French–American–British

Cooperative Study Group criteria were used for histological subgroup classification (Cheson et al. 1990). Randomly selected Bilkent University students, with no history of hematological malignancies or any other type of cancer ($n=124$), were genotyped in order for us to assess the status of the *MYH* mutations in apparently healthy Turkish individuals. Informed consent was obtained from all the students.

We screened for the *MYH* Tyr165Cys and Gly382Asp variants using genomic DNA as described (Al-Tassan et al. 2002). Neither mutation was present in any of the samples, except for that of one patient diagnosed with AML/M3. *MYH* Tyr165Cys mutation in the heterozygous state was present in the sample obtained at the time of initial diagnosis. Further sampling at remission, and the analysis of parental DNA, showed only the normal allele. Therefore, the mutation was considered to be specific for the leukemic blasts. It may be interesting to screen the whole *MYH* gene for mutations in hematological malignancies in the future, especially if increased transversions of G:C to T:A proved to be present in leukemic blasts. Based on these results, an association between childhood leukemias and the *MYH* missense variants Tyr165Cys and Gly382Asp was not observed. Also, these variants appear to be absent—if not at a very low frequency—in the Turkish population, contrary to the British population.

Acknowledgements This work was supported by SBAG-2513 from the Turkish Scientific and Technical Research Council (TÜBİTAK) and Bilkent University Research Fund.

References

- Al-Tassan N, Chmiel NH, Maynard J, Fleming N, Livingston AL, Williams GT, Hodges AK, Davies DR, David SS, Sampson JR, Cheadle JP (2002) Inherited variants of *MYH* associated with somatic G:C → T:A mutations in colorectal tumors. *Nat Genet* 30:227–232
- Cheson BD, Cassileth PA, Head DR, Schiffer CA, Bennett JM, Bloomfield CD, Brunning R, Gale RP, Grever MR, Keating MJ, Sawitsky A, Stass S, Weinstein H, Woods GW (1990) Report of the National Cancer Institute-sponsored workshop on definitions of diagnosis and response in acute myeloid leukemia. *J Clin Oncol* 8:813–819
- Guan Y, Manuel RC, Arvai AS, Parikh SS, Mol CD, Miller JH, Lloyd S, Tainer JA (1998) MutY catalytic core, mutant and bound adenine structures define specificity for DNA repair enzyme superfamily. *Nat Struct Biol* 5:1058–1064
- Horwitz M (1997) The genetics of familial leukemia. *Leukemia* 11:1347–1359
- Jones S, Emmerson P, Maynard J, Best JM, Jordan S, Williams GT, Sampson JR, Cheadle JP (2002) Biallelic germline mutations in *MYH* predispose to multiple colorectal adenoma and somatic G:C → T:A mutations. *Hum Mol Genet* 11:2961–2967
- Peltomäki P (2001) Deficient DNA mismatch repair: a common etiologic factor for colon cancer. *Hum Mol Genet* 10:735–740
- Ricciardone MD, Özçelik T, Cevher B, Özdağ H, Tuncer M, Gürgey A, Uzunalımoğlu Ö, Çetinkaya H, Tanyeli A, Erken E, Öztürk M (1999) Human *MLH1* deficiency predisposes to hematological malignancy and neurofibromatosis type 1. *Cancer Res* 59:290–293
- Whiteside D, McLeod R, Graham G, Steckley JL, Booth K, Somerville MJ, Andrew SE (2002) A homozygous germ-line mutation in the human *MSH2* gene predisposes to hematological malignancy and multiple café-au-lait spots. *Cancer Res* 62:359–362
- Wang Q, Lasset C, Desseigne F, Frappaz D, Bergeron C, Navarro C, Ruano E, Puisieux A (1999) Neurofibromatosis and early onset of cancers in *hMLH1*-deficient children. *Cancer Res* 59:294–297